

On Non-central Stirling Numbers of the First Kind

Milan Janjić

Abstract

It is shown in this note that non-central Stirling numbers $s(n, k, \alpha)$ of the first kind naturally appear in the expansion of derivatives of the function $x^{-\alpha} \ln^{\beta} x$, where α and β are arbitrary real numbers. We first obtain a recurrence relation for these numbers, and then, using Leibnitz rule we obtain an explicit formula for them. We also obtain a formula for $s(n, 1, \alpha)$ and then derive several combinatorial identities related to these numbers.

1 Introduction

We are dealing here with a special kind of numbers introduced by D. S. Mitrinović in his note [4]. In the paper [5] tables are given for the numbers which we called non-central Stirling numbers of the first kind. Following [3], they will be denoted by $s(n, k, \alpha)$. Several other names are in use for these numbers. One of them is r -Stirling numbers, as in [1]. The definition in this paper is restricted to the case when α is a nonnegative integer, and $\alpha \leq n$. L. Carlitz [2] used the name: weighted Stirling numbers. In the well known encyclopedia [6] they are called the generalized Stirling numbers. Here we use the name and the notation from the book [3]. For instance, $(\alpha)_n = \alpha(\alpha - 1) \cdots (\alpha - n + 1)$ are falling factorials, and $s(n, k)$ are Stirling numbers of the first kind.

2 Derivatives of $x^{-\alpha} \ln^{\beta} x$.

We shall investigate derivatives of the function

$$f(x) = x^{-\alpha} \ln^{\beta} x, \quad (\alpha, \beta \in \mathbb{R}),$$

obtaining them in two different way.

Theorem 1. *Let α be real, and n nonnegative integer. Then*

$$f^{(n)}(x) = x^{-\alpha-n} \sum_{i=0}^n s(n, i, \alpha) (\beta)_i \ln^{\beta-i} x. \quad (1)$$

where $s(n, i, \alpha)$, $(0 \leq i \leq n)$ are polynomials of α with integer coefficients.

Proof. The assertion is true for $n = 0$ if we take $s(0, 0, \alpha) = 1$.
Taking

$$s(1, 0, \alpha) = -\alpha, \quad s(1, 1, \alpha) = 1,$$

we see that the assertion is true for $n = 1$.

Suppose that the assertion is true for $n \geq 1$.

Taking derivative in (1) we obtain

$$\begin{aligned} f^{(n+1)}(x) &= (-\alpha - n)x^{-\alpha-n-1} \sum_{i=0}^n s(n, i, \alpha)(\beta)_i \ln^{\beta-i} x + \\ &\quad + x^{-\alpha-n-1} \sum_{i=0}^{n-1} s(n, i, \alpha)(\beta)_{i+1} \ln^{\beta-i-1} x = \\ &= x^{-\alpha-n-1} \left[(-\alpha - n) \sum_{i=0}^n s(n, i, \alpha)(\beta)_i \ln^{\beta-i} x + \sum_{i=0}^n s(n, i, \alpha)(\beta)_{i+1} \ln^{\beta-i-1} x \right] \end{aligned}$$

Replacing $i + 1$ by i in the second sum we obtain

$$\begin{aligned} f^{(n+1)}(x) &= x^{-\alpha-n-1} \left[\sum_{i=0}^n (-\alpha - n)s(n, i, \alpha)(\beta)_i + \sum_{i=1}^{n+1} s(n, i-1, \alpha)(\beta)_i \ln^{\beta-i} x \right] = \\ &= x^{\alpha-n-1} (\alpha - n)s(n, 0, \alpha) + \\ &\quad + x^{\alpha-n-1} \sum_{i=1}^n \left[(-\alpha - n)s(n, i, \alpha) + s(n, i-1, \alpha) \ln^{\beta-i} x \right] (\beta)_i + \\ &\quad + s(n, n, \alpha)(\beta)_{n+1} \ln^{\beta-n-1} x. \end{aligned}$$

It follows that the assertion is true if we take

$$\begin{aligned} s(n+1, 0, \alpha) &= (-\alpha - n)s(n, 0, \alpha), \\ s(n+1, i, \alpha) &= (-\alpha - n)s(n, i, \alpha) + s(n, i-1, \alpha), \quad (i = 1, \dots, n), \\ s(n+1, n+1, \alpha) &= s(n, n, \alpha). \end{aligned} \tag{2}$$

The preceding equation are well-known recurrence relations for non-central Stirling numbers of the first kind [3, pp.316].

Note 1. It is obvious that $s(n, i, 0) = s(n, i)$ are Stirling numbers of the first kind.

Since $p(0, 0, \alpha) = 1$, for from $n = 1, 2, \dots$ from the first equation in (2) we obtain:

$$p(n, 0, \alpha) = (-\alpha)_n,$$

and, since $p(1, 1, \alpha) = 1$, from the last equation in (2) follows:

$$p(n, n, \alpha) = 1, \quad (n = 0, 1, 2, \dots).$$

By the use of Leibnitz formula we shall obtain an explicit expression for $s(n, k, \alpha)$. The following equation holds:

$$f^{(n)}(x) = \sum_{k=0}^n \binom{n}{k} (x^\alpha)^{(k)} (\ln^\beta x)^{(n-k)}. \quad (3)$$

First we have

$$(x^{-\alpha})^{(k)} = (-\alpha)_k x^{\alpha-k}.$$

Using induction it is easy to prove that:

$$[f(\ln x)]^{(n)} = x^{-n} \sum_{k=1}^n s(n, k) f^{(k)}(t), \quad (t = \ln x).$$

Taking particular $f(t) = t^\beta$ we obtain:

$$(\ln^\beta x)^{(n-k)} = x^{-n+k} \sum_{i=1}^{n-k} s(n-k, i) (\beta)_i \ln^{\beta-i} x.$$

Replacing these in (3) we have the following.

Theorem 2. *Let $\alpha \neq 0$ be real, and $n, i, (i \leq n)$ be nonnegative integers. Then*

$$s(n, i, \alpha) = \sum_{k=0}^{n-i} \binom{n}{k} (-\alpha)_k s(n-k, i). \quad (4)$$

Note 2. *Theorem is true even in the case $\alpha = 0$ if we additionally define $(0)_0 = 1$.*

3 Some combinatorial identities

Taking $i = 1$ in (4) we obtain the following:

Corollary 1. *Let α be a real number, and n be a positive integer. Then*

$$s(n, 1, \alpha) = n! \sum_{k=0}^{n-1} (-1)^{n-k-1} \frac{\binom{-\alpha}{k}}{n-k}.$$

For $s(n, 1, \alpha)$ we have the following recurrence relation:

$$s(1, 1, \alpha) = 1, \quad s(n, 1, \alpha) = (-\alpha - n + 1)s(n-1, 1, \alpha) + (-\alpha)_{n-1}, \quad (n \geq 2). \quad (5)$$

Particular, we have $s(2, 1, \alpha) = -2\alpha - 1$.

We shall now prove that polynomials $r(n, 1, \alpha)$, $(n = 1, 2, \dots)$ defined by:

$$r(n, 1, \alpha) = \sum_{k=0}^{n-1} (k+1)s(n, k+1)(-\alpha)^k,$$

satisfy the above recurrence relation. For $n = 1$ it is obviously true.

Using two terms recurrence relations for Stirling numbers of the first kind we obtain:

$$\begin{aligned} r(n, 1, \alpha) &= \sum_{k=0}^{n-1} (k+1)[s(n-1, k) - (n-1)s(n-1, k+1)](-\alpha)^k = \\ &= \sum_{k=0}^{n-1} (k+1)s(n-1, k)(-\alpha)^k - (n-1) \sum_{k=0}^{n-2} (k+1)s(n-1, k+1)(-\alpha)^k. \end{aligned}$$

Since $s(n-1, 0) = 0$, by replacing $k+1$ instead of k in the first sum on the right we have:

$$\begin{aligned} r(n, -\alpha) &= \sum_{k=0}^{n-2} (k+2)s(n-1, k+1)(-\alpha)^{k+1} - (n-1) \sum_{k=0}^{n-2} (k+1)s(n-1, k+1)(-\alpha)^k = \\ &= (-\alpha - n + 1)r(n-1, \alpha) + \sum_{k=0}^{n-2} s(n-1, k+1)(-\alpha)^{k+1}. \end{aligned}$$

Furthermore, a well known property of Stirling numbers implies:

$$\sum_{k=0}^{n-2} s(n-1, k+1)(-\alpha)^{k+1} = \sum_{k=1}^{n-1} s(n-1, k)(-\alpha)^k = (-\alpha)_{n-1}.$$

We thus obtain that $r(n, 1, \alpha)$ satisfies (5). In this way we have proved the following identity:

Theorem 3. *Let α be a real number, and $n \geq 1$ be an integer. Then:*

$$n! \sum_{k=0}^{n-1} (-1)^k \frac{\binom{-\alpha}{k}}{n-k} = \sum_{k=0}^{n-1} (k+1)s(n, k+1)\alpha^k, \quad (6)$$

where $s(n, k+1)$ are unsigned Stirling numbers of the first kind.

Note 3. *Theorem is true for $\alpha = 0$ with the convention that $0^0 = 1$.*

Some particular values for α in (6) gives several interesting combinatorial identities. For $\alpha = -1$ we obtain an identity expressing factorials in terms of Stirling numbers of the first kind.

Corollary 2. *The following formula is true:*

$$(-1)^n (n-2)! = \sum_{k=0}^{n-1} (k+1)s(n, k+1), \quad (n \geq 2).$$

For $\alpha = 1$ we obtain a formula for the sum of reciprocals of natural numbers.

Corollary 3. *For each $n = 1, 2, \dots$ we have:*

$$n! \left(1 + \frac{1}{2} + \dots + \frac{1}{n} \right) = \sum_{k=0}^{n-1} (k+1) \mathbf{s}(n, k+1).$$

Consider now the case that $\alpha < 0$ is an integer and $n \geq 1 - \alpha$. In this case the function

$$q(n, 1, \alpha) = (-1)^{n+\alpha-1} (-\alpha)! (n + \alpha - 1)!, \quad (n \geq 1 - \alpha)$$

satisfies the recurrence relation (5). In fact, in this case we have $(-\alpha)_{n-1} = 0$, since a factor in this product must be zero. Thus:

$$\begin{aligned} (-\alpha - n + 1)q(n-1, 1, \alpha) &= -(n + \alpha - 1)(-1)^{n-\alpha-2} (-\alpha)! (n + \alpha - 2)! = \\ &= (-1)^{n-\alpha-1} (-\alpha)! (n + \alpha - 1)! = q(n, 1, \alpha). \end{aligned}$$

We have thus obtained the following:

Corollary 4. *Let n, α be positive integers and $n \geq \alpha + 1$. Then:*

$$n! \sum_{k=0}^{\alpha} \frac{(-1)^{\alpha-k} \binom{\alpha}{k}}{n-k} = \alpha! (n - \alpha - 1)!.$$

In other words, the following equation holds:

$$(\alpha + 1) \sum_{k=0}^{\alpha} \frac{(-1)^{\alpha-k} \binom{\alpha}{k}}{n-k} = \frac{1}{\binom{n}{\alpha+1}}.$$

In the case that α is a negative integer and $n \leq -\alpha$ the identity (6) is closely related to the harmonic numbers. Namely, denote $H_0 = 0$, $H_n = \sum_{k=1}^n \frac{1}{k}$, ($n = 1, 2, \dots$), and define:

$$h(n, 1, \alpha) = (H_{-\alpha} - H_{-\alpha-n}) \frac{(-\alpha)!}{(-\alpha-n)!}, \quad (\alpha = -1, -2, \dots; n = 1, 2, \dots, -\alpha).$$

It will be shown that $h(n, 1, \alpha)$ satisfies (5). We use the induction on n . For $n = 1$ we have:

$$h(1, 1, \alpha) = \frac{(-\alpha)!}{(-\alpha-1)!} (H_{-\alpha} - H_{-\alpha-1}) = 1.$$

Furthermore we have:

$$\begin{aligned} (-\alpha - n + 1)h(n-1, 1, \alpha) + (-\alpha)_{n-1} &= \frac{(-\alpha)!}{(-\alpha-n)!} (H_{-\alpha} - H_{-\alpha-n+1}) + (-\alpha)_{n-1} = \\ &= \frac{(-\alpha)!}{(-\alpha-n)!} (H_{-\alpha} - H_{-\alpha-n}) - \frac{(-\alpha)!}{(-\alpha-n)!(-\alpha-n+1)} + (-\alpha)_{n-1} = \end{aligned}$$

$$= \frac{(-\alpha)!}{(-\alpha-n)!} (H_{-\alpha} - H_{-\alpha-n}),$$

since $\frac{(-\alpha)!}{(-\alpha-n)!(-\alpha-n+1)!} = (-\alpha)_{n-1}$. We thus obtain:

$$h(n, 1, \alpha) = (-\alpha - n + 1)b(n - 1, 1, \alpha) + (-\alpha)_{n-1},$$

that is, $h(n, 1, \alpha)$ satisfies (5). We have thus proved the following.

Corollary 5. *Let α be a positive integer and $1 \leq n \leq \alpha$ be integers. Then:*

$$H_{\alpha} - H_{\alpha-n} = \frac{(-1)^{n+1}}{\binom{\alpha}{n}} \sum_{k=0}^{n-1} \frac{(-1)^k \binom{\alpha}{k}}{n-k}.$$

Equivalently:

$$H_{\alpha} - H_{\alpha-n} = \frac{\sum_{k=0}^{n-1} (k+1)s(n, k+1)\alpha^k}{\sum_{k=0}^n s(n, k)\alpha^k}.$$

In the case $\alpha = n$ we have the following expressions for harmonic numbers.

Corollary 6. *Let n be a positive integer. Then:*

$$H_n = (-1)^{n+1} \sum_{k=0}^{n-1} \frac{(-1)^k \binom{n}{k}}{n-k},$$

and

$$H_n = \frac{1}{n!} \sum_{k=0}^{n-1} (k+1)s(n, k+1)n^k.$$

References

- [1] A. Z. Broder, The r -Stirling numbers, it Department of Computer Science, Stanford University, Stanford, CA 94305, 1982.
- [2] L. Carlitz, Weighted Stirling Numbers of the First and Second kind. IT. *The Fibonacci Quarterly* 18(1980):242-257.
- [3] Ch. A. Charalambides, Enumerative Combinatorics, *Chapman & Hall/CRC*, Boca Raton, Florida, 2002.
- [4] D. S. Mitrinović, Sur une classe de nombres relies aux nombres de Stirling. *Les Comptes rendus de l'Academie des Sciences de Paris*, 1. 252, 1961, p. 2354-2356.
- [5] D. S. Mitrinović et R. S. Mitrinović, Tableaux d'une classe de nombres relies aux nombres de Stirling. *Univ. Beograd. Pubi. Elektrotehn. Fak. Ser. Mat. Fiz. No. 77* 1962, 77 pp.
- [6] N. J. Sloane, The Encyclopedia of Integer Sequences. Electronically published at www.research.att.com